

PROVIDER DATA SHEET

Female Fertility Profiles I and II



The Problem

In America, a large number of couples are having difficulty trying to get pregnant. Current statistics from the most recent National Survey of Family Growth¹ show that 7.4% (2.1 million) of all married women of childbearing age (15-44 years) in the US are infertile, defined as not having achieved pregnancy despite using no contraception in the past 12 months or more. When the survey considered only married, childless women, this figure increases to 16.6%. The incidence of female infertility is significantly age-related, being higher with increasing age. In one population study², the main causes of infertility attributable to the woman were ovulatory failure (21%), tubal damage (14%), and endometriosis (6%), while a massive 28% of cases were unexplained. Yet in the absence of a physical cause, many cases of female infertility may be explained by something as simple as a correctable hormone imbalance, which can be assessed by hormone testing. This involves the collection of saliva and a simple finger-stick to obtain drops of blood that are deposited on a filter paper and allowed to dry.

Hormones tested in the Female Fertility Profiles

Hormones tested in both profiles are: estradiol (E2), progesterone (Pg), testosterone (T), DHEA-S, diurnal cortisol (C) sampled four times during a day, thyroid hormones (free T3, free T4, TSH, and TPO antibodies), LH and FSH. There are two profiles to select from: Profile I is saliva dominant, meaning that the sex (E2, Pg, T) and adrenal (DHEA-S, C) hormones are tested in saliva, while the thyroid hormones, LH and FSH are tested in blood spot. For those who prefer to have the sex hormones tested in blood instead of saliva, we created Profile II, which tests all of the hormones in blood spot with the exception of the 4x diurnal C tested in saliva. Sampling is done on days 19-21 of the menstrual cycle, coinciding with the mid-luteal phase, when the level of progesterone should be optimal for a successful pregnancy.

Hormonal Aspects of Infertility

Hormone-related causes of female infertility most often involve the following five scenarios:

1. Ovarian Insufficiency

While the average age of menopause is 51, women can start experiencing signs of ovarian insufficiency even in their thirties³. A cessation of ovulation prior to the age of 40 is rare, and is usually referred to as premature ovarian failure. Declining ovarian function is the main reason for the age-related decline in female fertility. As the number of available follicles starts to fall, estrogen is still being produced but ovulation does not occur, and progesterone levels fall in the absence of a corpus luteum⁴. Also, while high FSH levels on day 3 of the menstrual cycle typically confirm premature ovarian failure and the onset of menopause, FSH levels remain elevated on day 19-21 compared to those seen in normal ovulatory cycles, and despite normal LH values^{5,6}. A typical pattern of day 19-21 hormone levels indicating signs of ovarian insufficiency would consist of: low estradiol, low progesterone, low testosterone and elevated FSH; LH, DHEA-S, cortisol and thyroid hormones may or may not be normal.

2. Luteal Phase Deficiency

In some patients who are infertile, ovulation may occur normally but levels of progesterone are inadequate during the luteal phase. This luteal progesterone deficiency means that, even if the egg is fertilized, implantation either does not occur, or if it does the progesterone produced by the corpus luteum is not high enough to sustain the pregnancy. Luteal phase deficiency can be caused by a number of problems, including endometriosis⁷ and abnormal follicular development, but most commonly it is a result of inadequate progesterone production by the corpus luteum. A typical finding is low progesterone levels in the luteal phase, usually with normal estradiol levels⁸.

3. Polycystic Ovarian Syndrome (PCOS)

PCOS is the most common endocrine disorder affecting women of reproductive age and is closely associated with insulin resistance, metabolic syndrome and future risk of developing diabetes and cardiovascular disease⁹. Among women presenting
(over)

with infertility in one study, PCOS was found to be present in 81% of women who were anovulatory, in 50% of those with tubal disease, and in 44% of those with unexplained infertility¹⁰. PCOS is diagnosed when a patient has two of these three criteria: polycystic ovaries on ultrasound; oligo- and/or anovulation; and hyperandrogenism⁹. Hormonally, it is characterized by low progesterone, normal-to-high estradiol, high testosterone, normal to high DHEA-S, and LH elevated 2-3 times relative to FSH. Cortisol and thyroid hormones may or may not be normal, although women with PCOS have been found to have a three-fold higher prevalence of autoimmune thyroiditis compared to healthy women¹¹.

4. Hypometabolism/Thyroid Deficiency

Thyroid dysfunction, including subclinical hypothyroidism (elevated TSH with normal free T3 and free T4 levels), has been implicated as a cause of infertility, yet thyroid hormone treatment can be a simple solution to restore a regular menstrual pattern¹². In one study, levothyroxine treatment resulted in pregnancy in 44% of infertile patients diagnosed with subclinical hypothyroidism¹³. In patients with hypometabolism as a result of thyroid dysfunction, the sex hormones, LH and FSH and cortisol may all be normal but the patient has symptoms of hypothyroidism, and a low T3 level. High TPO antibodies indicate an autoimmune thyroid disease (e.g., Hashimoto's Disease), which is associated with fertility-related problems, and it is important to rule out thyroid autoimmunity in women attempting to conceive because of the increased risk of miscarriage¹⁴.

5. Stress

Stress can severely affect a woman's ability to conceive¹⁵, probably because of its impact on endocrine balance, particularly affecting thyroid and adrenal function and the pituitary's production of gonadotrophs. (See www.endotext.com¹⁶ for a comprehensive, illustrated overview of the effects of stress on endocrine physiology). The diurnal cortisol variation measured in saliva samples collected on waking, late morning, late afternoon and at bedtime, indicate the effects of stress on adrenal function as well as the adrenal glands' ability to produce cortisol, a hormone important for the cellular actions of thyroid hormones. Stress stimuli also increase prolactin production, and high cortisol and prolactin levels are seen in endometriosis. Endometriosis is found in more than 50% of women with unexplained infertility, and the high cortisol and prolactin levels induced by stress have been implicated in the development of this condition¹⁷.

Advantages of a Simple Combined Saliva/ Blood Spot Test for Infertility

This simple, cost-effective test can identify many problems related to hormonal imbalances associated with infertility.

- Convenient, private sample collection can be done at home and mailed in for analysis
- Excellent correlation of blood spot and plasma/serum levels of analytes¹⁸
- Hormone results can indicate a correctable hormonal imbalance

Useful websites:

International Council on Infertility Information Dissemination: www.inciid.org
American Society for Reproductive Medicine: www.asrm.org
American Academy of Fertility Care Professionals: www.aafcp.org
NaProTechnology (Natural Procreative Technology): www.naprotechnology.com

References

1. Chandra A, Martinez GM, Mosher WD, Abma JC, Jones J. Fertility, Family Planning, and Reproductive Health of US Women: Data from the 2002 National Survey of Family Growth. National Center for Health Statistics. *Vital Health Stat* 2005; 23(25). Available at: http://www.cdc.gov/nchs/data/series/sr_23/sr23_025.pdf (accessed 3/17/2008).
2. Hull MG, Glazener CM, Kelly NJ, Conway DL, Foster PA, Hinton RA, Coulson C, Lambert PA, Watt EM, Desai KM. Population study of causes, treatment, and outcome of infertility. *Br Med J (Clin Res Ed)*. 1985 Dec 14;291(6510):1693-7.
3. National Institute on Aging, US National Institutes of Health. The Age Page. Available at: <http://www.niapublications.org/agepages/menopause.asp> (accessed 3/17/2008).
4. Ahmed Ebbiary NA, Lenton EA, Salt C, Ward AM, Cooke ID. The significance of elevated basal follicle stimulating hormone in regularly menstruating infertile women. *Hum Reprod*. 1994;9(2):245-52.
5. Reame NE, Kelche RP, Beitins IZ, Yu MY, Zawacki CM, Padmanabhan V. Age effects of follicle-stimulating hormone and pulsatile luteinizing hormone secretion across the menstrual cycle of premenopausal women. *J Clin Endocrinol Metab*. 1996;81(4):1512-8.
6. Santoro N, Brown JR, Adel T, Skurnick JH. Characterization of reproductive hormonal dynamics in the perimenopause. *J Clin Endocrinol Metab*. 1996;81(4):1495-501.
7. Cunha-Filho JS, Gross JL, Bastos de Souza CA, Lemos NA, Giugliani C, Freitas F, Passos EP. Physiopathological aspects of corpus luteum defect in infertile patients with mild/minimal endometriosis. *J Assist Reprod Genet*. 2003;20(3):117-21.
8. Soules MR, McLachlan RI, Ek M, Dahl KD, Cohen NL, Bremner WJ. Luteal phase deficiency: characterization of reproductive hormones over the menstrual cycle. *J Clin Endocrinol Metab*. 1989;69(4):804-12.
9. Kousta E, Tolis G, Franks S. Polycystic ovary syndrome. Revised diagnostic criteria and long-term health consequences. *Hormones (Athens)*. 2005;4(3):133-47.
10. Kousta E, White DM, Cela E, McCarthy MI, Franks S. The prevalence of polycystic ovaries in women with infertility. *Hum Reprod*. 1999;14(11):2720-3.
11. Janssen OE, Mehlmauer N, Hahn S, Offner AH, Gärtner R. High prevalence of autoimmune thyroiditis in patients with polycystic ovary syndrome. *Eur J Endocrinol*. 2004;150(3):363-9.
12. Trokoudes KM, Skordis N, Picolos MK. Infertility and thyroid disorders. *Curr Opin Obstet Gynecol*. 2006;18(4):446-51.
13. Abalovich M, Mitelberg L, Allami C, Gutierrez S, Alcaraz G, Otero P, Levalle O. Subclinical hypothyroidism and thyroid autoimmunity in women with infertility. *Gynecol Endocrinol*. 2007; 23(5):279-83.
14. Poppe K, Glinoeer D, Tournaye H, Devroey P, Schiettecatte J, Haentjens P, Velkeniers B. Thyroid autoimmunity and female infertility. *Verh K Acad Geneesk Belg*. 2006;68(5-6):357-77.
15. Nakamura K, Sheps S, Clara Arck P. Stress and reproductive failure: past notions, present insights and future directions. *J Assist Reprod Genet*. 2008;25(2-3):47-62.
16. Tsigos K, Cyrou I, Chrousos G. Stress, endocrine physiology and pathophysiology. Available at: <http://www.endotext.com/adrenal/adrenal8/adrenalframe8.htm> (accessed 4/7/08).
17. Lima AP, Moura MD, Rosa e Silva AA. Prolactin and cortisol levels in women with endometriosis. *Braz J Med Biol Res*. 2006;39(8):1121-7.
18. Edelman A, Stouffer R, Zava DT, Jensen JT. A comparison of blood spot vs. plasma analysis of gonadotropin and ovarian steroid hormone levels in reproductive-age women. *Fertil Steril*. 2007;88(5):1404-7.